ГОЛИШНИК НА СОФИЙСКИЯ УНИВЕРСИТЕТ "СВ. КЛИМЕНТ ОХРИДСКИ" ФАКУЛТЕТ ПО МАТЕМАТИКА И ИНФОРМАТИКА Том 98

ANNUAIRE DE L'UNIVERSITE DE SOFIA "ST. KLIMENT OHRIDSKI"
FACULTE DE MATHEMATIQUES ET INFORMATIQUE
Tome 98

AN EXAMPLE OF A 16-VERTEX FOLKMAN EDGE (3,4)-GRAPH WITHOUT 8-CLIQUES

NIKOLAY KOLEV, NEDYALKO NENOV

In [6] we computed the edge Folkman number F(3,4;8) = 16. There we used and announced without proof that in any blue-red coloring of the edges of the graph $K_1 + C_5 + C_5$ there is either a blue 3-clique or red 4-clique. In this paper we give a detailed proof of this fact.

Keywords: Folkman graph, Folkman number

2000 MSC: 05C55

1. INTRODUCTION

Only finite non-oriented graphs without multiple edges and loops are considered. We call a p-clique of the graph G a set of p vertices each two of which are adjacent. The largest positive integer p such that G contains a p-clique is denoted by $\operatorname{cl}(G)$. A set of vertices of the graph G none two of which are adjacent is called an independent set. In this paper we shall also use the following notations:

- V(G) is the vertex set of the graph G;
- E(G) is the edge set of the graph G;
- N(v), $v \in V(G)$ is the set of all vertices of G adjacent to v;
- G[V], $V \subseteq V(G)$ is the subgraph of G induced by V;
- $\chi(G)$ is the chromatic number of G;

- K_n is the complete graph on n vertices;
- C_n is the simple cycle on n vertices.

The equality $C_n = v_1 v_2 \dots v_n$ means that $V(C_n) = \{v_1, \dots, v_n\}$ and

$$E(C_n) = \{[v_i, v_{i+1}], i = 1, \dots, n-1\} \cup \{[v_1, v_n]\}$$

Let G_1 and G_2 be two graphs without common vertices. We denote by $G_1 + G_2$ the graph G for which $V(G) = V(G_1) \cup V(G_2)$ and $E(G) = E(G_1) \cup E(G_2) \cup E'$ where $E' = \{[x, y] : x \in V(G_1), y \in V(G_2)\}.$

Let G and H be two graphs. We shall say that H is a subgraph of G and we shall denote $H \subseteq G$ when $V(H) \subseteq V(G)$ and $E(H) \subseteq E(G)$.

Definition 1.1. A 2-coloring

$$E(G) = E_1 \cup E_2, \qquad E_1 \cap E_2 = \emptyset, \tag{1.1}$$

is called a blue-red coloring of the edges of the graph G (the edges in E_1 are blue and the edges in E_2 are red).

We define for blue-red coloring (1.1) and for an arbitrary vertex $v \in V(G)$

$$N_i(v) = \{x \in N(v) \mid [v, x] \in E_i\}, \quad i = 1, 2;$$

 $G_i(v) = G[N_i(v)].$

Definition 1.2. Let H be a subgraph of G. We say that H is a monochromatic subgraph in the blue-red coloring (1.1) if $E(H) \subseteq E_1$ or $E(H) \subseteq E_2$. If $E(H) \subseteq E_1$ we say that H is a blue subgraph, and if $E(H) \subseteq E_2$ we say that H is a red subgraph.

Definition 1.3. The blue-red coloring (1.1) is called (p,q)-free, if there are no blue p-cliques and no red q-cliques. The symbol $G \to (p,q)$ means that any blue-red coloring of E(G) is not (p,q)-free. If $G \to (p,q)$ then G is called edge Folkman (p,q)-graph.

Let p, q and r be positive integers. The Folkman number F(p, q; r) is defined by the equality

$$F(p,q;r) = \min\{|V(G)| : G \to (p,q) \text{ and } \operatorname{cl}(G) < r\}.$$

In [1] Folkman proved that

$$F(p,q;r)$$
 exists $\iff r > \max\{p,q\}.$

That is why the numbers F(p,q;r) are called Folkman numbers. Only few Folkman numbers are known. An exposition of the results on the Folkman numbers was given in [6]. In [6] we computed a new Folkman number, namely F(3,4;8) = 16. This result is based upon the fact that $K_1 + C_5 + C_5 + C_5 \rightarrow (3,4)$, which was announced without proof in [6]. In this paper we give a detailed proof of this fact. So, the aim of this paper is to prove the following

Main Theorem. Let $G = K_1 + C_5^{(1)} + C_5^{(2)} + C_5^{(3)}$, where $C_5^{(1)}$, $C_5^{(2)}$, $C_5^{(3)}$ are copies of the 5-cycle C_5 . Then $G \to (3,4)$.

2. AUXILIARY RESULTS

Lemma 2.1. Let $E(G) = E_1 \cup E_2$ be a (3,4)-free red-blue coloring of the edges of the graph G. Then:

- (a) $G_1(v)$ is a red subgraph, $v \in V(G)$;
- (b) $(E(G_2(v)) \cap E_1) \cup (E(G_2(v)) \cap E_2)$ is a (3,3)-free red-blue coloring of $E(G_2(v))$, $v \in V(G)$. Thus $G_2(v) \not\rightarrow (3,3)$.

Proof. The statement of (a) is obvious. Assume that (b) is not true. Then, since there is no blue 3-clique, $G_2(v)$ contains a red 3-clique. This red 3-clique together with the vertex v form a red 4-clique, which is a contradiction. \square

Corollary 2.1. Let $E(G) = E_1 \cup E_2$ be a (3,4)-free blue-red coloring of E(G). Then:

- (a) $cl(G_1(v)) \leq 3, v \in V(G);$
- (b) $cl(G_2(v)) \le 5, v \in V(G);$
- (c) $G_2(v) \not\supseteq K_3 + C_5, v \in V(G)$.

Proof. The statement of (a) follows from Lemma 2.1(a). The statements of (b) and (c) follow from Lemma 2.1(b), since $K_6 \to (3,3)$, [4] and $K_3 + C_5 \to (3,3)$, [2]. \square

Lemma 2.2 ([5]). Let $G = C_5 + H$, where $V(H) = \{x, y, z\}$ and $E(H) = \{[x, y], [x, z]\}$. Let $E(G) = E_1 \cup E_2$ be a (3, 3)-free blue-red coloring of E(G). Then H is monochromatic in this coloring.

Lemma 2.3 ([3]). Let $G = C_5 + K_2$ and $E(G) = E_1 \cup E_2$ be a (3,3)-free blue-red coloring of E(G) such that $E(C_5) \subseteq E_i$. Then $E(K_2) \in E_i$.

Lemma 2.4. Let $G = K_1 + C_5^{(1)} + C_5^{(2)} + C_5^{(3)}$, where $C_5^{(1)}$, $C_5^{(2)}$, $C_5^{(3)}$ are copies of the 5-cycle C_5 and $V(K_1) = \{a\}$. Let $E(G) = E_1 \cup E_2$ be a blue-red coloring of E(G) such that $cl(G_1(a)) \leq 3$ and $G_2(a) \neq (3,3)$. Then, up to numeration of the 5-cycles $C_5^{(1)}$, $C_5^{(2)}$ and $C_5^{(3)}$, we have:

- (a) $N_1(a) \supset V(C_5^{(1)})$ and $N_1(a) \cap V(C_5^{(2)})$ is an independent set;
- (b) $N_2(a) \supset V(C_5^{(3)})$ and $N_2(a) \cap V(C_5^{(2)})$ is not an independent set.

Proof. Let $C_5^{(1)} = v_1 v_2 v_3 v_4 v_5$, $C_5^{(2)} = u_1 u_2 u_3 u_4 u_5$ and $C_5^{(3)} = w_1 w_2 w_3 w_4 w_5$. We shall use the following obvious fact

$$\chi(C_5) = 3. \tag{2.1}$$

It follows from (2.1) that

$$N_1(a) \cap V(C_5^{(i)})$$
 or $N_2(a) \cap V(C_5^{(i)})$ is not an independent set, $i=1,2,3$. (2.2)

By (2.2) and Corollary 2.1(b), at least one of the sets $N_2(a) \cap V(C_5^{(i)})$, i = 1, 2, 3, is an independent set. Thus, at least one of the sets $N_1(a) \cap V(C_5^{(i)})$, i = 1, 2, 3, is not an independent set. Without loss of generality we can assume that

$$N_1(a) \cap V(C_5^{(1)})$$
 is not an independent set. (2.3)

It follows from Corollary 2.1(a) and (2.3) that $N_1(a) \cap V(C_5^{(2)}) = \emptyset$ or $N_1(a) \cap V(C_5^{(3)}) = \emptyset$. Let for example $N_1(a) \cap V(C_5^{(3)}) = \emptyset$. Then

$$N_2(a) \supset V(C_5^{(3)}).$$
 (2.4)

We have from (2.3) and Corollary 2.1(a) that $N_1(a) \cap V(C_5^{(2)})$ is an independent set. Thus, it follows from (2.1) that $N_2(a) \cap V(C_5^{(2)})$ is not an independent set. This fact, together with (2.4) and Corollary 2.1(c), gives $N_2(a) \cap V(C_5^{(1)}) = \emptyset$. Hence, $N_1(a) \supseteq V(C_5^{(1)})$. The Lemma is proved. \square

Lemma 2.5. Let $G = K_1 + C_5^{(1)} + C_5^{(2)} + C_5^{(3)}$, where $C_5^{(i)}$, i = 1, 2, 3, are copies of the 5-cycle C_5 . Let $E(G) = E_1 \cup E_2$ be a blue-red coloring such that some of the cycles $C_5^{(1)}$, $C_5^{(2)}$, $C_5^{(3)}$ is not monochromatic. Then this coloring is not (3,4)-free.

Proof. Let $V(K_1) = \{a\}$, $C_5^{(1)} = v_1v_2v_3v_4v_5$, $C_5^{(2)} = u_1u_2u_3u_4u_5$ and $C_5^{(3)} = w_1w_2w_3w_4w_5$. Assume the opposite, i.e. $E(G) = E_1 \cup E_2$ is (3,4)-free. Then by Corollary 2.1(a) we have $\operatorname{cl}(G_1(a)) \leq 3$ and by Lemma 2.1(b) we have $G_2(a) \not\to (3,3)$. Thus, according to Lemma 2.4 we can assume that

$$N_1(a) \supseteq V(C_5^{(1)}) \text{ and } N_1(a) \cap V(C_5^{(2)}) \text{ is independent;}$$
 (2.5)

$$N_2(a) \supseteq V(C_5^{(3)})$$
 and $N_2(a) \cap V(C_5^{(2)})$ is not independent. (2.6)

It follows from (2.5) and Lemma 2.1(a) that

$$E(C_5^{(1)}) \subseteq E_2. \tag{2.7}$$

We have from the statement of the Lemma 2.5 that at least one of the cycles $C_5^{(i)}$, i = 1, 2, 3, is not monochromatic and since $E(C_5^{(1)}) \subseteq E_2$ it remains to consider the following two cases:

Case 1. $C_5^{(2)}$ is not monochromatic. Let for example $[u_1, u_5] \in E_1$ and $[u_1, u_2] \in E_2$. If $u_1, u_2, u_5 \in N_2(a)$ by (2.6) we have $G_2(a) \supset C_2^{(3)} + G[u_1, u_2, u_5]$. It follows from Lemma 2.2 that $G_2(a)$ contains a monochromatic 3-clique. This contradicts Lemma 2.1(b). So, at least one of the vertices u_1, u_2, u_5 belongs to $N_1(a)$. Therefore, we have the following subcases:

Subcase 1a. $u_1 \in N_1(a)$. Since there are no blue 3-cliques it follows from (2.5) that

$$N_2(u_1) \supset V(C_5^{(1)}).$$
 (2.8)

As $[u_1, a]$, $[u_1, u_5] \in E_1$ and $cl(G_1(u_1)) \leq 3$ (see Corollary 2.1(a)), the set $N_1(u_1) \cap V(C_5^{(3)})$ is independent. Therefore, $N_2(u_1) \cap V(C_5^{(3)})$ is not independent. This fact, together with $[u_1, u_2] \in E_2$ and (2.8), gives $G_2(u_1) \supset K_3 + C_5^{(1)}$, which contradicts Corollary 2.1(c).

Subcase 1b. $u_2 \in N_1(a)$ and $u_1 \in N_2(a)$. Since there are no blue 3-cliques it follows from (2.5) that

$$N_2(u_2) \supset V(C_5^{(1)}).$$
 (2.9)

If $N_2(u_1) \cap V(C_5^{(1)})$ contains two adjacent vertices then these vertices together with u_1 and u_2 form a red 4-clique according to (2.7) and (2.9). Hence $N_2(u_1) \cap V(C_5^{(1)})$ is independent and, therefore, $N_1(u_1) \cap V(C_5^{(1)})$ is not independent. Since $u_5 \in N_1(u_1)$ and $cl(G_1(u_1)) \leq 3$ (see Corollary 2.1(a)) we have $N_1(u_1) \cap V(C_5^{(3)}) = \emptyset$. Hence

$$N_2(u_1) \supset V(C_5^{(3)}).$$
 (2.10)

By (2.6) and (2.10)

$$V(C_5^{(3)}) \subseteq N_2(u_1) \cap N_2(a)$$
.

Since $[a, u_1] \in E_2$ and there are no red 4-cliques we obtain that

$$E(C_5^{(3)}) \subseteq E_1.$$
 (2.11)

As there are no blue 3-cliques from (2.11) it follows that $N_1(u_2) \cap V(C_5^{(3)})$ is independent. Therefore, $N_2(u_1) \cap V(C_5^{(3)})$ contains two adjacent vertices. This fact, together with $[u_1, u_2] \in E_2$ and (2.9), gives $G_2(u_2) \supset K_3 + C_5^{(1)}$, which contradicts Corollary 2.1(c).

Subcase 1c. $u_5 \in N_1(a)$ and $u_1, u_2 \in N_2(a)$. Since $a, u_2 \in N_2(u_1)$, it follows from Corollary 2.1(b) that at least one of the sets $N_2(u_1) \cap V(C_5^{(1)})$ and $N_2(u_1) \cap V(C_5^{(3)})$ is independent. Hence at least one of the sets $N_1(u_1) \cap V(C_5^{(1)})$, $N_1(u_1) \cap V(C_5^{(3)})$, is not independent. Assume that $N_1(u_1) \cap V(C_5^{(1)})$ is not independent. This fact, together with $u_5 \in N_1(u_1)$ and Corollary 2.1(a), implies

$$N_2(u_1) \supset V(C_5^{(3)}).$$
 (2.12)

As $[a, u_1, u_2]$ is a red 3-clique and $[a, u_1, u_2, w_i]$ is not a red 4-clique, $i = 1, \ldots, 5$, it follows from (2.6) and (2.12) that $[u_2, w_i] \in E_1$, $i = 1, \ldots, 5$, i.e. $N_1(u_2) \supset$

 $V(C_5^{(3)})$. We have from Lemma 2.1(a) that $E(C_5^{(3)}) \subseteq E_2$. Thus, according to (2.6) and (2.12), the vertices a and u_1 together with two adjacent vertices of $C_5^{(3)}$ form a red 4-clique, which is a contradiction.

Let us now consider the situation when $N_1(u_1) \cap V(C_5^{(3)})$ is not independent. Corollary 2.1(a) and $u_5 \in N_1(u_1)$ imply

$$N_2(u_1) \supset V(C_5^{(1)}).$$
 (2.13)

If $N_2(u_1) \cap V(C_5^{(3)}) \neq \emptyset$ then from $a, u_2 \in N_2(u_1)$ and (2.13) it follows that $G_2(u_1) \supset K_3 + C_5^{(1)}$, which contradicts the Corollary 2.1(c). Hence $N_2(u_1) \cap V(C_5^{(3)}) = \emptyset$, i.e.

$$N_1(u_1) \supset C_5^{(3)}$$
. (2.14)

Since there are no blue 3-cliques, we obtain from (2.14) and Lemma 2.1(a):

$$E(C_5^{(3)}) \subseteq E_2.$$
 (2.15)

If $N_2(u_2) \cap V(C_5^{(3)})$ is not independent then according to (2.6) and (2.15) an edge in $N_2(u_2) \cap V(C_5^{(3)})$ together with a and u_2 form a red 4-clique. Let $N_2(u_2) \cap V(C_5^{(3)})$ be independent. Then $N_1(u_2) \cap V(C_5^{(3)})$ is not independent. Thus, it follows from Corollary 2.1(a) that $N_1(u_2) \cap V(C_5^{(1)})$ is independent and $N_2(u_2) \cap V(C_5^{(1)})$ is not independent. Then an edge in $N_2(u_2) \cap V(C_5^{(1)})$, together with the vertices u_1 and u_2 , form a red 4-clique, according to (2.7) and (2.13), which is a contradiction.

Case 2. $C_5^{(3)}$ is not monochromatic but $C_5^{(2)}$ is monochromatic. Without loss of generality we can assume that $[w_1, w_5] \in E_1$ and $[w_1, w_2] \in E_2$. Since $a, w_2 \in N_2(w_1)$ it follows from Corollary 2.1(b) that at least one of the sets $N_2(w_1) \cap V(C_5^{(1)})$ and $N_2(w_1) \cap V(C_5^{(2)})$ is independent. Hence at least one of the sets $N_1(w_1) \cap V(C_5^{(1)})$, $N_1(w_1) \cap V(C_5^{(2)})$ is not independent. We shall consider these possibilities:

Subcase 2a. $N_1(w_1) \cap V(C_5^{(1)})$ is not independent. Since $[w_1, w_5] \in E_1$ it follows from Corollary 2.1(a) that $N_1(w_1) \cap V(C_5^{(2)}) = \emptyset$, i.e.

$$N_2(w_1) \supset V(C_5^{(2)}).$$
 (2.16)

By Lemma 2.1(b) $G_2(w_1)$ does not contain a monochromatic 3-clique and $G_2(w_1) \supset C_5^{(2)} + [a, w_2]$. Since $C_5^{(2)}$ is monochromatic and $[a, w_2] \in E_2$, it follows from Lemma 2.3 that

$$E(C_5^{(2)}) \subseteq E_2.$$
 (2.17)

We see from (2.6), (2.16) and (2.17) that the vertices a and w_1 , together with an edge of $C_5^{(2)}$ form a red 4-clique, which is a contradiction.

Subcase 2b. $N_1(w_1) \cap V(C_5^{(2)})$ is not independent. Since $w_5 \in N_1(w_1)$ it follows from Corollary 2.1(a) that

$$N_2(w_1) \supset V(C_5^{(1)}).$$
 (2.18)

Corollary 2.1(c) and $G_2(w_1) \supset C_5^{(1)} + [a, w_2] = K_2 + C_5^{(1)}$ imply

$$N_1(w_1) \supset V(C_5^{(2)}).$$
 (2.19)

Lemma 2.1(a) and (2.19) give

$$E(C_5^{(2)}) \subseteq E_2.$$
 (2.20)

Since there are no blue 3-cliques and $[w_1, w_5] \in E_1$ it follows from (2.19) that

$$N_2(w_5) \supset V(C_5^{(2)}).$$
 (2.21)

We see from (2.6), (2.20) and (2.21) that the vertices a and w_5 together with an edge of $C_5^{(2)}$ form a red 4-clique which is a contradiction. \square

3. A PROPERTY OF THE GRAPH $C_5 + C_5 + C_5$

Let $G = C_5^{(1)} + C_5^{(2)} + C_5^{(3)}$ where $C_5^{(i)}$, i = 1, 2, 3, are copies of the 5-cycle C_5 . Let us consider the blue-red coloring where $E_1 = E(C_5^{(1)}) \cup E(C_5^{(2)}) \cup E(C_5^{(3)})$. It is clear that this coloring is (3,4)-free. Thus $G \not\to (3,4)$. However the following theorem holds:

Theorem 3.1. Let $G = C_5^{(1)} + C_5^{(2)} + C_5^{(3)}$ where $C_5^{(i)}$, i = 1, 2, 3, are copies of the 5-cycle C_5 . Let $E(G) = E_1 \cup E_2$ be a blue-red coloring such that $E(C_5^{(1)}) \subseteq E_2$, $E(C_5^{(2)}) \subseteq E_1$ and $E(C_5^{(3)}) \subseteq E_1$. Then this coloring is not (3, 4)-free.

Proof. Assume the opposite, i.e. that there are no blue 3-cliques and no red 4-cliques. Let $C_5^{(1)} = v_1 v_2 v_3 v_4 v_5$, $C_5^{(2)} = u_1 u_2 u_3 u_4 u_5$, $C_5^{(3)} = w_1 w_2 w_3 w_4 w_5$. Since the cycles $C_5^{(2)}$ and $C_5^{(3)}$ are blue and there are no blue 3-cliques, the sets $N_1(v_i) \cap V(C_5^{(2)})$ and $N_1(v_i) \cap V(C_5^{(3)})$ are independent. Thus, we have

$$|N_2(v_i) \cap V(C_5^{(2)})| \ge 3, \quad |N_2(v_i) \cap V(C_5^{(3)})| \ge 3, \quad i = 1, \dots, 5.$$
 (3.1)

It follows from (3.1) that

$$N_2(x) \cap N_2(y) \cap V(C_5^{(i)}) \neq \emptyset, \quad i = 2, 3, \quad x, y \in V(C_5^{(1)}).$$
 (3.2)

Let $x, y \in V(C_5^{(1)})$. We define

$$B_1(x,y) = \{ v \in V(C_5^{(2)}) \mid [x,v], [y,v] \in E_2 \},$$

$$B_2(x,y) = \{ v \in V(C_5^{(3)}) \mid [x,v], [y,v] \in E_2 \}.$$

We see from (3.2) that

$$B_i(x,y) \neq \emptyset, \quad i = 1, 2, \quad x, y \in V(C_5^{(1)}).$$
 (3.3)

Ann. Sofia Univ., Fac. Math. and Inf., 98, 2008, 127-141.

We shall prove that

if
$$[x, y] \in E(C_5^{(1)})$$
 then $B_i(x, y)$ is independent, $i = 1, 2$. (3.4)

Assume the opposite and let for example $u', u'' \in B_1(x, y)$ and $[u', u''] \in E(C_5^{(2)})$. By (3.3) there exists $w \in B_2(x, y)$. Since there are no blue 3-cliques, then at least one of the edges [u', w], [u'', w] is red. Hence [x, y, u', w] or [x, y, u'', w] is a red 4-clique, which is a contradiction.

Let u' and u'' be adjacent vertices in $C_5^{(2)}$. Since $[u', u''] \in E_1$ and there are no blue 3-cliques, we have

$$N_1(u') \cap N_1(u'') \cap V(C_5^{(1)}) = \emptyset.$$

Thus $|N_1(u') \cap V(C_5^{(1)})| \le 2$ or $|N_1(u'') \cap V(C_5^{(1)})| \le 2$. Hence

$$|N_2(u') \cap V(C_5^{(1)})| \ge 3 \text{ or } |N_2(u'') \cap V(C_5^{(1)})| \ge 3.$$
 (3.5)

So, (3.5) holds for every two adjacent vertices in $C_5^{(2)}$. Hence $|N_2(u) \cap V(C_5^{(1)})| \geq 3$ holds for at least three vertices in $C_5^{(2)}$. Thus, there exist two adjacent vertices in $C_5^{(2)}$, for example u_1 and u_2 , such that

$$|N_2(u_1) \cap V(C_5^{(1)})| \ge 3 \text{ and } |N_2(u_2) \cap V(C_5^{(1)})| \ge 3.$$
 (3.6)

If the both inequalities in (3.6) are strict then $N_2(u_1) \cap N_2(u_2) \cap V(C_5^{(1)})$ contains two adjacent vertices v' and v''. Since $u_1, u_2 \in B(v', v'')$ then this contradicts (3.4). Thus, we may assume that $|N_2(u_1) \cap V(C_5^{(1)})| = 3$. Hence $N_2(u_1) \cap V(C_5^{(1)})$ contains two adjacent vertices, for example v_3 and v_4 . Now we shall prove that the third vertex in $N_2(u_1) \cap V(C_5^{(1)})$ is the vertex v_1 . Assume the opposite. Then $v_2 \in N_2(u_1) \cap V(C_5^{(1)})$ or $v_5 \in N_2(u_1) \cap V(C_5^{(1)})$. Let $v_2 \in N_2(u_1) \cap V(C_5^{(1)})$. Then $v_1, v_5 \in N_1(u_1)$. Since $v_1, v_5, u_2 \in N_1(u_1)$ it follows from Corollary 2.1(a) that $N_1(u_1) \cap V(C_5^{(3)}) = \emptyset$. Thus, $G_2(u_1)$ contains $C_5^{(3)} + [v_3, v_4] = K_2 + C_5$. According to Lemma 2.1(b) $G_2(u_1)$ does not contain monochromatic 3-cliques. As $E(C_5^{(3)}) \subseteq E_1$ and $[v_3, v_4] \in E_2$, this contradicts Lemma 2.3. We proved that $v_2 \notin N_2(u_1)$. Analogously we prove that $v_5 \notin N_2(u_1)$. So,

$$v_1, v_3, v_4 \in N_2(u_1) \text{ and } v_2, v_5 \in N_1(u_1).$$
 (3.7)

By (3.3) we can assume that $w_1 \in B_2(v_3, v_4)$. Since $[v_3, v_4, u_1, w_1]$ is not a red 4-clique, we have

$$[u_1, w_1] \in E_1. \tag{3.8}$$

As there are no blue 3-cliques and $[u_1, v_2]$, $[u_1, v_5] \in E_1$, it follows that $[w_1, v_2]$, $[w_1, v_5] \in E_2$. Taking into consideration $w_1 \in B_2(v_3, v_4)$, we have

$$[w_1, v_i] \in E_2, \quad i = 2, 3, 4, 5.$$
 (3.9)

By (3.3) there is $u \in B_1(v_2, v_3)$. Since $[v_2, u_1] \in E_1$ then $u \neq u_1$. We shall prove that $u = u_3$ or $u = u_4$. Assume the opposite. Then $u = u_2$ or $u = u_5$. Let, for example, $u = u_2$. Since $[v_2, v_3, u_2, w_1]$ is not a red 4-clique, it follows from (3.9) and $u_2 \in B_1(v_2, v_3)$ that $[u_2, w_1] \in E_1$. We obtained the blue 3-clique $[u_1, u_2, w_1]$, which is a contradiction. This contradiction proves that $u = u_3$ or $u = u_4$. We can assume without loss of generality that $u = u_3$. We have

$$[u_3, w_1] \in E_1, \tag{3.10}$$

because $[v_2, v_3, u_3, w_1]$ is not a red 4-clique. By (3.3) there exists $u \in B_1(v_4, v_5)$. Repeating the above considerations about $u \in B_1(v_2, v_3)$ we see that $u = u_3$ or $u = u_4$.

Case 1. $u = u_4$. Since $[v_4, v_5, w_1, u_4]$ is not a red 4-clique, we have $[u_4, w_1] \in E_1$. Hence $[u_3, u_4, w_1]$ is a blue 3-clique, which is a contradiction.

Case 2. $u = u_3$. In this case we have $u_3 \in B_1(v_2, v_3) \cap B_1(v_4, v_5)$, i.e.

$$[u_3, v_i] \in E_2, \quad i = 2, 3, 4, 5.$$
 (3.11)

As $[v_1, w_1, u_3]$ is not a blue 3-clique, it follows from (3.10) that $[v_1, u_3] \in E_2$ or $[v_1, w_1] \in E_2$.

Subcase 2a. $[v_1, u_3] \in E_2$. By (3.11) $N_2(u_3) \supset C_5^{(1)}$. Since there are no blue 3-cliques $N_2(u_3)$ contains two adjacent vertices $w', w'' \in V(C_5^{(3)})$. Thus $G_2(u_3) \supset C_5^{(1)} + [w', w'']$. By Lemma 2.1(b) $G_2(u_3)$ contains no monochromatic 3-cliques. This contradicts Lemma 2.3 because $E(C_5^{(1)}) \subseteq E_2$ and $[w', w''] \in E_1$.

Subcase 2b. $[v_1, w_1] \in E_2$. By (3.9) we see that $N_2(w_1) \supset V(C_5^{(1)})$. Since there are no blue 3-cliques $N_2(w_1)$ contains two adjacent vertices $u', u'' \in V(C_5^{(2)})$. Hence $N_2(w_1) \supset C_5^{(1)} + [u', u'']$ which contradicts Lemma 2.3.

The theorem is proved. \square

4. PROOF OF THE MAIN THEOREM

Let $C_5^{(1)} = v_1 v_2 v_3 v_4 v_5$, $C_5^{(2)} = u_1 u_2 u_3 u_4 u_5$, $C_5^{(3)} = w_1 w_2 w_3 w_4 w_5$ and $V(K_1) = \{a\}$. Assume the opposite, i.e. there exists a (3,4)-free blue-red coloring $E_1 \cup E_2$ of the edges of $K_1 + C_5^{(1)} + C_5^{(2)} + C_5^{(3)}$. By Lemma 2.4 we can assume that:

$$N_1(a) \supset V(C_5^{(1)})$$
 and $N_1(a) \cap V(C_5^{(2)})$ is independent; (4.1)

$$N_2(a) \supset V(C_5^{(3)})$$
 and $N_2(a) \cap V(C_5^{(2)})$ is not independent. (4.2)

We shall prove that

$$E(C_5^{(i)}) \subseteq E_2, \quad i = 1, 2, 3.$$
 (4.3)

By (4.1) and Lemma 2.1(a), $E(C_5^{(1)}) \subseteq E_2$. According to Lemma 2.5 each of the 5-cycles $C_5^{(2)}$ and $C_5^{(3)}$ is monochromatic. By (4.2) $G_2(a) \supset C_5^{(3)} + e$ where $e \in E(C_5^{(2)})$. By Lemma 2.1(b) $G_2(a)$ contains no monochromatic 3-cliques. Thus, it follows from Lemma 2.3 that the edge e and the 5-cycle $C_5^{(3)}$ have the same color. Therefore, the 5-cycles $C_5^{(2)}$ and $C_5^{(3)}$ are monochromatic of the same color. Thus, it follows from Theorem 3.1 that $E(C_5^{(2)}) \nsubseteq E_1$ and $E(C_5^{(3)}) \nsubseteq E_1$. We proved (4.3). Now we shall prove that

$$N_2(a) = V(C_5^{(2)}) \cup V(C_5^{(3)}). \tag{4.4}$$

Assume the opposite. Then it follows from (4.2) that $N_1(a) \cap V(C_5^{(2)}) \neq \emptyset$. Let for example $u_1 \in N_1(a) \cap V(C_5^{(2)})$, i.e. $[u_1, a] \in E_1$. We see from (4.1) that

$$[a, u_2] \in E_2. \tag{4.5}$$

As there are no blue 3-cliques by (4.1) and $[u_1, a] \in E_1$ we obtain

$$N_2(u_1) \supset V(C_5^{(1)}).$$
 (4.6)

We see from Corollary 2.1(a) that at least one of the sets $N_2(u_2) \cap V(C_5^{(3)})$, $N_2(u_2) \cap V(C_5^{(1)})$ is not independent. If $N_2(u_2) \cap V(C_5^{(1)})$ is not independent, it follows from (4.6) and (4.3) that the vertices u_1 and u_2 together with an edge of $C_5^{(1)}$ form a red 4-clique. If $N_2(u_2) \cap V(C_5^{(3)})$ is not independent, by (4.3), (4.5) and (4.2), the vertices a and u_2 together with an edge of $C_5^{(3)}$ form a red 4-clique. This contradiction proves (4.4).

It follows from (4.4) and Lemma 2.1(b) that

$$C_5^{(2)} + C_5^{(3)}$$
 contains no monochromatic 3-cliques. (4.7)

Now we obtain from (4.7) and (4.3)

$$N_2(x) \cap V(C_5^{(3)})$$
 is independent, $x \in V(C_5^{(2)});$ (4.8)

$$N_2(x) \cap V(C_5^{(2)})$$
 is independent, $x \in V(C_5^{(3)})$. (4.9)

Let us note that

$$N_1(x) \cap V(C_5^{(1)})$$
 is independent, $x \in V(C_5^{(2)}) \cup V(C_5^{(3)})$. (4.10)

Indeed, let for example $x \in V(C_5^{(2)})$. By (4.8), $N_1(x) \cap V(C_5^{(3)})$ is not independent. This fact and Corollary 2.1(a) prove (4.10).

We shall prove that

$$N_1(x) \cap V(C_5^{(2)}), \ x \in V(C_5^{(1)}) \text{ is not independent } \iff N_2(x) \supset V(C_5^{(3)}); \ (4.11)$$

$$N_1(x) \cap V(C_5^{(3)}), \ x \in V(C_5^{(1)})$$
 is not independent $\iff N_2(x) \supset V(C_5^{(2)}).$ (4.12)

The statements (4.11) and (4.12) are proved analogously. That is why we shall prove (4.11) only. Let $N_1(x) \cap V(C_5^{(2)})$, $x \in V(C_5^{(1)})$ be not independent. Since $[x,a] \in E_1$, it follows from Corollary 2.1(a) that $N_1(x) \cap V(C_5^{(3)}) = \emptyset$, i.e. $N_2(x) \supset V(C_5^{(3)})$. Now let $N_2(x) \supset V(C_5^{(3)})$, $x \in V(C_5^{(1)})$. Assume that $N_1(x) \cap V(C_5^{(2)})$ is independent. Then $N_2(x) \cap V(C_5^{(2)})$ is not independent. Since $C_5^{(1)}$ is red, $G_2(x) \supset K_3 + C_5^{(3)}$ which contradicts Corollary 2.1(c). So, (4.11) and (4.12) are proved. Using (4.11) and (4.12) we shall prove that

$$N_1(x) \cap V(C_5^{(i)}), i = 2, 3, \text{ is independent}, x \in V(C_5^{(1)}).$$
 (4.13)

Assume that (4.13) is wrong and let, for example, $N_1(v_1) \cap V(C_5^{(2)})$ be not independent (remind that $C_5^{(1)} = v_1v_2v_3v_4v_5$). Then by (4.11) $N_2(v_1) \supset V(C_5^{(3)})$. If $N_2(v_2) \cap V(C_5^{(3)})$ is not independent then v_1 and v_2 together with two adjacent vertices from $N_2(v_2) \cap V(C_5^{(3)})$ form a red 4-clique, which is a contradiction. Therefore, $N_1(v_2) \cap V(C_5^{(3)})$ is not independent. Thus (4.12) gives $N_2(v_2) \supset V(C_5^{(2)})$. Repeating the above considerations about the vertex v_1 on v_2 we obtain $N_2(v_3) \supset V(C_5^{(3)})$. In the same way it follows from $N_2(v_3) \supset V(C_5^{(3)})$ that $N_2(v_4) \supset V(C_5^{(2)})$. At the end it follows from $N_2(v_4) \supset V(C_5^{(2)})$ that $N_2(v_5) \supset V(C_5^{(3)})$. So, we proved that

$$N_2(v_1) \cap N_2(v_5) \supset V(C_5^{(3)}).$$

Thus, it follows from (4.3) that v_1 and v_5 , together with an edge of $C_5^{(3)}$, form a red 4-clique, which is a contradiction. This contradiction proves (4.13). According to (4.13) it follows from (4.11) and (4.12) that

$$N_2(x) \not\supset V(C_5^{(i)}), \ i = 2, 3, \quad x \in V(C_5^{(1)}).$$
 (4.14)

Let $x \in V(C_5^{(2)}) \cup V(C_5^{(3)})$. By (4.10) $|N_1(x) \cap V(C_5^{(1)})| \le 2$. Thus, we have the following possibilities:

Case 1. $N_1(x) \cap V(C_5^{(1)}) = \emptyset$ for some vertex $x \in V(C_5^{(2)}) \cup V(C_5^{(3)})$. Let, for example, $N_1(u_1) \cap V(C_5^{(1)}) = \emptyset$ (remind that $C_5^{(2)} = u_1u_2u_3u_4u_5$). Then $N_2(u_1) \supset V(C_5^{(1)})$. We see from (4.10) that $N_2(u_2) \cap V(C_5^{(1)})$ is not independent. Thus u_1 and u_2 , together with two adjacent vertices from $N_2(u_2) \cap V(C_5^{(1)})$, form a red 4-clique, which is a contradiction.

Case 2. $|N_1(x) \cap V(C_5^{(1)})| = 1$ for some vertex $x \in V(C_5^{(2)}) \cup V(C_5^{(3)})$. Let, for example, $|N_1(u_1) \cap V(C_5^{(1)})| = 1$. Without loss of generality we can consider $[u_1, v_1] \in E_1$ and $[u_1, v_i] \in E_2$, i = 2, 3, 4, 5. According to (4.14) we can assume that $[v_1, w_1] \in E_1$. Since there are no blue 3-cliques, $[u_1, w_1] \in E_2$. It follows from (4.10) that $N_2(w_1) \cap V(C_5^{(1)})$ contains two adjacent vertices. As

$$N_2(w_1) \cap V(C_5^{(1)}) \subseteq N_2(u_1) \cap V(C_5^{(1)}) = \{v_2, v_3, v_4, v_5\}$$

we see that u_1 and w_1 together with two adjacent vertices in $\{v_2, v_3, v_4, v_5\}$ form a red 4-clique, which is a contradiction.

Case 3. $|N_1(x) \cap V(C_5^{(1)})| = 2$ for every $x \in V(C_5^{(2)}) \cup V(C_5^{(3)})$. According to (4.8) $N_1(u_1) \cap V(C_5^{(3)})$ is not independent. Thus, we can assume that $w_1, w_2 \in N_1(u_1) \cap V(C_5^{(3)})$, i.e.

$$[u_1, w_1], [u_1, w_2] \in E_1.$$
 (4.15)

It follows from (4.13)

$$N_1(w_1) \cap N_1(w_2) \cap V(C_5^{(1)}) = \emptyset.$$
 (4.16)

In the case considered we have

$$|N_1(w_1) \cap V(C_5^{(1)})| = |N_1(w_2) \cap V(C_5^{(1)})| = |N_1(u_1) \cap V(C_5^{(1)})| = 2.$$

We obtain from (4.16)

$$N_1(u_1) \cap N_1(w_1) \cap V(C_5^{(1)}) \neq \emptyset \text{ or } N_1(u_1) \cap N_1(w_2) \cap V(C_5^{(1)}) \neq \emptyset.$$

By (4.15) there is a blue 3-clique, which is a contradiction. The Main Theorem is proved.

5. EXAMPLE OF FOLKMAN EDGE (3,5)-GRAPH WITHOUT 13-CLIQUES

Using the Main Theorem we shall prove the following

Theorem 5.1. Let $G = K_4 + C_5^{(1)} + C_5^{(2)} + C_5^{(3)} + C_5^{(4)}$ where $C_5^{(i)}$, i = 1, ..., 4, are copies of the 5-cycle C_5 . Then $G \to (3,5)$.

In order to prove Theorem 5.1 we shall need the following

Lemma 5.1. Let $E(G) = E_1 \cup E_2$ is a (3,5)-free blue-red coloring of E(G). Then:

- (a) $G_1(v)$, $v \in V(G)$, is a red subgraph;
- (b) $(E(G_2(v)) \cap E_1) \cup (E(G_2(v)) \cap E_2)$ is a (3,4)-free blue-red coloring of $E(G_2(v))$, $v \in V(G)$. Thus, $G_2(v) \not\to (3,4)$.

Lemma 5.1 is proved in the same way as Lemma 2.1.

Corollary 5.1. Let $E(G) = E_1 \cup E_2$ be a (3,5)-free blue-red coloring of E(G). Then:

- (a) $cl(G_1(v)) \le 4, v \in V(G)$;
- (b) $cl(G_2(v)) \le 8, v \in V(G);$

- (c) $G_2(v) \not\supset K_4 + C_5 + C_5$;
- (d) $G_2(v) \not\supset K_1 + C_5 + C_5 + C_5$.

Proof. The statement (a) follows from Lemma 5.1(a). The statement (b) follows from Lemma 5.1(b) and $K_9 \to (3,4)$, [4]. The statement (c) follows from Lemma 5.1(b) and $K_4 + C_5 + C_5 \to (3,4)$, [8]. The statement (d) follows from Lemma 5.1(b) and the Main Theorem. \square

Proof of Theorem 5.1. Assume the opposite, i.e. there exists a blue-red coloring $E(G) = E_1 \cup E_2$, which is (3,5)-free. Let $V(K_4) = \{a_1, a_2, a_3, a_4\}$.

Case 1. There exists $a_i \in V(K_4)$ such that $|N_1(a_i) \cap V(K_4)| = 3$. Let, for example, $[a_1, a_2]$, $[a_1, a_3]$, $[a_1, a_4] \in E_1$. By Corollary 5.1(a) at most one of the sets $N_1(a_1) \cap V(C_5^{(i)})$, i = 1, 2, 3, 4, is not empty, i.e. $N_2(a_1)$ contains at least three of the cycles $C_5^{(i)}$, i = 1, 2, 3, 4. Let, for example,

$$N_2(a_1) \supset V(C_5^{(2)}) \cup V(C_5^{(3)}) \cup V(C_5^{(4)}).$$

By Corollary 5.1(a) it follows that $N_1(a_1) \cap V(C_5^{(1)})$ is independent. Thus, $N_2(a_1) \cap V(C_5^{(1)}) \neq \emptyset$. We obtained that $G_2(a_1) \supset K_1 + C_5^{(2)} + C_5^{(3)} + C_5^{(4)}$, which contradicts Corollary 5.1(d).

Case 2. There exists $a_i \in V(K_4)$ such that $|N_1(a_i) \cap V(K_4)| = 2$. Let, for example, $[a_1, a_2]$, $[a_1, a_3] \in E_1$ and $[a_1, a_4] \in E_2$. Since $[a_1, a_4] \in E_2$ if the sets $N_2(a_1) \cap V(C_5^{(i)})$, i = 1, 2, 3, 4, are not independent then $G_2(a_1) \supset K_9$, which contradicts Corollary 5.1(b). Hence, at least one of the sets $N_1(a_1) \cap V(C_5^{(i)})$, i = 1, 2, 3, 4, is not independent. Let, for example, $N_1(a_1) \cap V(C_5^{(1)})$ be not independent. According to Corollary 5.1(a) it follows from this fact and $[a_1, a_2]$, $[a_1, a_3] \in E_1$ that $N_1(a_1) \cap V(C_5^{(i)})\emptyset$, i = 2, 3, 4, i.e. $N_2(a_1) \supset V(C_5^{(i)})$, i = 2, 3, 4. As $[a_1, a_4] \in E_2$ we have $G_2(a_1) \supset K_1 + C_5^{(2)} + C_5^{(3)} + C_5^{(4)}$, which contradicts Corollary 5.1(d).

Case 3. There exist $a_i \in V(K_4)$ such that $|N_1(a_i) \cap V(K_4)| = 1$. Let, for example, $[a_1, a_2] \in E_1$ and $[a_1, a_3]$, $[a_1, a_4] \in E_2$. We see from Corollary 5.1(a) that at least three of the sets $N_2(a_1) \cap V(C_5^{(i)})$, i = 1, 2, 3, 4, are not independent. Let, for example, $N_2(a_1) \cap V(C_5^{(2)})$, $N_2(a_1) \cap V(C_5^{(3)})$ and $N_2(a_1) \cap V(C_5^{(4)})$ be not independent. Since $[a_1, a_3]$, $[a_1, a_4] \in E_2$ it follows from Corollary 5.1(b) that $N_1(a_1) \supset V(C_5^{(1)})$. According to Lemma 5.1(a) it follows from this fact and $[a_1, a_2] \in E_1$ that at least two of the sets $N_1(a_1) \cap V(C_5^{(i)})$, i = 2, 3, 4, are empty. Therefore, we can assume that $N_2(a_1) \supset V(C_5^{(3)})$ and $N_2(a_1) \supset V(C_5^{(4)})$. Since $N_2(a_1) \cap V(C_5^{(2)})$ is not independent we have $G_2(a_1) \supset K_4 + V(C_5^{(3)}) + V(C_5^{(4)})$, which contradicts Corollary 5.1(c).

Case 4. $E(K_4) \subseteq E_2$. Since $[a_1, a_i] \in E_2$, i = 2, 3, 4, it follows from Corollary 5.1(b) that at least two of the sets $N_1(a_1) \cap V(C_5^{(i)})$, i = 1, 2, 3, 4, are not

independent. Let, for example, $N_1(a_1) \cap V(C_5^{(1)})$ and $N_1(a_1) \cap V(C_5^{(2)})$ be not independent. Then by Corollary 5.1(a) $N_1(a_1) \cap V(C_5^{(3)}) = \emptyset$ and $N_1(a_1) \cap V(C_5^{(4)}) = \emptyset$, i.e.

$$N_2(a_1) \supseteq V(C_5^{(3)}) \cup V(C_5^{(4)}).$$
 (5.1)

Since $[a_1, a_i] \in E_2$, i = 2, 3, 4, it follows from (5.1) and Corollary 5.1(c) that $N_2(a_1) \cap V(C_5^{(1)}) = \emptyset$ and $N_2(a_1) \cap V(C_5^{(2)}) = \emptyset$. That is why, we have from (5.1)

$$N_1(a_1) = V(C_5^{(1)}) \cup V(C_5^{(2)}). \tag{5.2}$$

As the vertices a_1 , a_2 , a_3 , a_4 are equivalent, in this case the above considerations prove that

$$N_1(a_i)$$
, $i = 1, 2, 3, 4$, is a union of two of the cycles $C_5^{(1)}$, $C_5^{(2)}$, $C_5^{(3)}$, $C_5^{(4)}$. (5.3)

Lemma 5.1(a) and (5.2) imply

$$C_5^{(1)} + C_5^{(2)}$$
 is a red subgraph. (5.4)

Since there are no red 5-cliques, we see from (5.4) that

$$N_1(a_i) \cap V(C_5^{(1)}) \neq \emptyset$$
 or $N_1(a_i) \cap V(C_5^{(2)}) \neq \emptyset$, $i = 2, 3, 4$.

Thus, by (5.3) we have

$$N_1(a_i) \supset V(C_5^{(1)}) \text{ or } N_1(a_i) \supset V(C_5^{(2)}), \quad i = 2, 3, 4.$$
 (5.5)

Hence, we can assume that

$$N_1(a_2) \supset V(C_5^{(1)})$$
 and $N_1(a_3) \supset V(C_5^{(1)})$. (5.6)

Let $C_5^{(1)} = v_1 v_2 v_3 v_4 v_5$. By (5.3) we have the following possibilities:

Subcase 4a. $N_1(a_4) \supset V(C_5^{(1)})$. According to (5.6) and (5.2) $[v_1, a_i] \in E_1$, i = 1, 2, 3, 4. Hence, by Corollary 5.1(a) $N_1(v_1) \cap V(C_5^{(i)}) = \emptyset$, i = 2, 3, 4, i.e. $G_2(v_1) \supset C_5^{(2)} + C_5^{(3)} + C_5^{(4)}$. By (5.2) $[v_1, v_2] \in E_2$. Thus, $G_2(v_1) \supset K_1 + C_5^{(2)} + C_5^{(3)} + C_5^{(4)}$, which contradicts Corollary 5.1(d).

Subcase 4b. $N_1(a_4) \cap V(C_5^{(1)}) = \emptyset$, i.e. $N_2(a_4) \supset V(C_5^{(1)})$. We have, from (5.2) and (5.6), $[v_1, a_i] \in E_1$, i = 1, 2, 3, and $[v_1, a_4] \in E_2$. By Corollary 5.1(a), at least two of the sets $N_1(v_1) \cap V(C_5^{(i)})$, i = 2, 3, 4, are empty. Thus, we can assume that

$$G_2(v_1) \supset C_5^{(3)} + C_5^{(4)}.$$
 (5.7)

It follows from Corollary 5.1(a) that $N_1(v_1) \cap V(C_5^{(2)})$ is independent. Hence, $N_2(v_1) \cap V(C_5^{(2)})$ is not independent. This fact, together with $[v_1, v_2], [v_1, a_4] \in E_2$ and (5.7), gives $G_2(v_1) \supset K_4 + C_5^{(3)} + C_5^{(4)}$, which contradicts Corollary 5.1(c). This contradiction finishes the proof of Theorem 5.1. \square

Since cl(G) = 12 and |V(G)| = 24, Theorem 5.1 implies

Corollary 5.2. $F(3,5;13) \le 24$.

Lin proved in [7] that $F(3,5;13) \ge 18$. In [9] Nenov improved this result, proving that either $K_8 + C_5 + C_5 \rightarrow (3,5)$ or $F(3,5;13) \ge 19$.

REFERENCES

- Folkman, J. Graphs with monochromatic complete subgraphs in every edge coloring. SIAM J. Appl. Math., 18, 1970, 19–24.
- Graham, R. On edgewise 2-colored graph with monohromatic triangles containing no complete hexagon. J. Combin. Theory, 4, 1968, 300.
- Graham, R., J. Spencer. On small graphs with forced monochromatic triangles. Recent Trends in Graph Theory, Lecture Notes in Math, 186, 1971, 137–141.
- Greenwood, R., A. Gleason. Combinatorial relation and chromatic graphs. Canad. J. Math., 7, 1955, 1–7.
- Khadzhiivanov, N., N. Nenov. On the Graham-Spencer number. Compt. rend. Acad. bulg. Sci., 31, 1978, 631-633.
- 6. Kolev, N., N. Nenov. The Folkman number $F_c(3,4;8)$ is equal to 16. Compt. rend. Acad. bulg. Sci., **59**, 2006, No 1, 25–30.
- 7. Lin, S. On Ramsey number and K_r -coloring of graphs. J. Comb. Theory, **B12**, 1972, 82–92.
- Nenov, N. On the Ramsey (3, 4)-graphs. Ann. Sofia Univ., Fac. Math. Mech., 73, 1979, 185–190 (1986) (in Russian).
- 9. Nenov, N. On the Zykov numbers and some its applications to Ramsey theory. Serdica Bulg. Math. Publ., 9, 1983, 161–167 (in Russian).

Received on November 15, 2005

Faculty of Mathematics and Informatics "St. Kl. Ohridski" University of Sofia 5, J. Bourchier blvd., 1164 Sofia BULGARIA

E-mail: nenov@fmi.uni-sofia.bg

E-mail: nickyxy@mail.bg